

THE POINCARÉ SERIES OF THE COVARIANTS OF BINARY FORMS

L. BEDRATYUK

ABSTRACT. A formula for computation of the Poincaré series $P_d(z)$ of the algebra of the covariants of binary d -form is found. By using it, we have computed the $P_d(z)$ for $d \leq 20$.

1. Consider the natural action of the complex matrix group SL_2 on the complex vector space of binary d -forms. Let C_d be corresponding algebra of covariants. The algebra C_d is a finite generated graded algebra:

$$C_d = (C_d)_0 + (C_d)_1 + \cdots + (C_d)_i + \cdots,$$

and the vector spaces $(C_d)_i$ are all finite dimensional. The formal power series $P_d(z) \in \mathbb{Z}[[z]]$,

$$P_d(z) = \sum_{i=0}^{\infty} \dim((C_d)_i) z^i,$$

is called the Poincaré series of the algebra of covariants C_d . The finite generation of the algebra of covariants implies that its Poincaré series is an expansion of some rational function. We consider here the problem of computing efficiently this rational function.

The Poincaré series calculations were an important object of research in classical invariant theory of the 19th century. For the cases $d \leq 10$, $d = 12$ the series $P_d(z)$ were calculated by Sylvester and Franklin, see [1], [2]. Relatively recently, Springer [3] set the explicit formula for computing the Poincaré series of the algebras of invariants of the binary d -forms. This formula has been used by Brouwer and Cohen [4] for $d \leq 17$ and also by Littelmann and Procesi [5] for even $d \leq 36$.

In the paper we have found a Sylvester-Cayley-type formula for calculating of $\dim(C_d)_i$ and Springer-type formula for calculation of $P_d(z)$. By using the formula, the series $P_d(z)$ is calculated for $d \leq 30$.

2. To begin with, we give a proof of the Sylvester-Cayley-type formula for covariants of binary form.

Let $V \cong \mathbb{C}^2$ be standard two-dimensional representation of Lie algebra $\mathfrak{sl}_2$. The irreducible representation $V_d = \langle v_0, v_1, \dots, v_d \rangle$, $\dim V_d = d + 1$ of the algebra $\mathfrak{sl}_2$ is the symmetric d -power of the standard representation $V = V_1$, i.e. $V_d = S^d(V)$, $V_0 \cong \mathbb{C}$. The basis elements $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ of the algebra $\mathfrak{sl}_2$ act on V_d by the derivations D_1, D_2, E :

$$D_1(v_i) = i v_{i-1}, D_2(v_i) = (d - i) v_{i+1}, E(v_i) = (d - 2i) v_i.$$

The action of $\mathfrak{sl}_2$ is extended to action on the symmetrical algebra $S(V_d)$ in the natural way. Let $\mathfrak{u}_2$ be the maximal unipotent subalgebra of $\mathfrak{sl}_2$. The algebra $\mathcal{S}_d$

$$\mathcal{S}_d := S(V_d)^{\mathfrak{u}_2} = \{v \in S(V_d) | D_1(v) = 0\},$$

is called the algebra of *seminvariants* of the binary form of degree d . For any element $v \in \mathcal{S}_d$ a natural number s is called the *order* of the element v if the number s is the smallest natural

number such that

$$D_2^s(v) \neq 0, D_2^{s+1}(v) = 0.$$

It is clear that any seminvariant $v \in S_d$ of order i is the highest weight vector for an irreducible $\mathfrak{sl}_2$ -module of the dimension $i + 1$ in $S(V_d)$.

The classical Robert's theorem [6] implies an isomorphism of the algebras covariants and seminvariants. Thus, it is enough to compute the Poincaré series of the algebra S_d .

The algebra $S(V_d)$ is graded

$$S(V_d) = S^0(V_d) + S^1(V_d) + \cdots + S^n(V_d) + \cdots,$$

and each $S^n(V_d)$ is the complete reducibly representation of the Lie algebra $\mathfrak{sl}_2$. Thus, the following decomposition holds

$$S^n(V_d) \cong \gamma_d(n, 0)V_0 + \gamma_d(n, 1)V_1 + \cdots + \gamma_d(n, dn)V_{dn}, \quad (*)$$

here $\gamma_d(n, k)$ is the multiplicity of the representation V_k in the decomposition of $S^n(V_d)$. On the other hand, the multiplicity $\gamma_d(n, i)$ of the representation V_i is equal to the number of linearly independent homogeneous seminvariants of degree n and order i for the binary d -form. This argument proves

Lema 1.

$$\dim(C_d)_n = \gamma_d(n, 0) + \gamma_d(n, 1) + \cdots + \gamma_d(n, dn).$$

The set of weights (eigenvalues of the operator E) of a representation W denote by Λ_W , in particular, $\Lambda_{V_d} = \{-d, -d+2, \dots, d\}$.

A formal sum

$$\text{Char}(W) = \sum_{k \in \Lambda_W} n_W(k)q^k,$$

is called the character of a representation W , here $n_W(k)$ denotes the multiplicity of the weight $k \in \Lambda_W$. Since, a multiplicity of any weight of the irreducible representation V_d is equal to 1, we have

$$\text{Char}(V_d) = q^{-d} + q^{-d+2} + \cdots + q^d.$$

The character $\text{Char}(S^n(V_d))$ of the representation $S^n(V_d)$ equals $H_n(q^{-d}, q^{-d+2}, \dots, q^d)$, see [7], where $H_n(x_0, x_1, \dots, x_d)$ is the complete symmetrical function

$$H_n(x_0, x_1, \dots, x_d) = \sum_{|\alpha|=n} x_0^{\alpha_0} x_1^{\alpha_1} \cdots x_d^{\alpha_d}, |\alpha| = \sum_i \alpha_i.$$

By replacing x_k with q^{d-2k} , $k = 0, \dots, d$, we obtain the specialized expression for the character of $\text{Char}(S^n(V_d))$:

$$\begin{aligned} \text{Char}(S^n(V_d)) &= \sum_{|\alpha|=n} (q^d)^{\alpha_0} (q^{d-2 \cdot 1})^{\alpha_1} \cdots (q^{d-2d})^{\alpha_d} = \\ &= \sum_{|\alpha|=n} q^{dn-2(\alpha_1+2\alpha_2+\cdots+d\alpha_d)} = \sum_{k=0}^{dn} \omega_d(n, k) q^{dn-2k}, \end{aligned}$$

here $\omega_d(n, k)$ is the number nonnegative integer solutions of the equation

$$\alpha_1 + 2\alpha_2 + \cdots + d\alpha_d = \frac{dn - k}{2}$$

on the assumption that $\alpha_0 + \alpha_1 + \cdots + d\alpha_d = n$. In particular, the coefficient of q^0 (the multiplicity of zero weight) is equal to $\omega_d(n, \frac{dn}{2})$, and the coefficient of q^1 is equal $\omega_d(n, \frac{dn-1}{2})$.

On the other hand, the decomposition $(*)$ implies the equality for the characters:

$$\text{Char}(S^n(V_d)) = \gamma_d(n, 0)\text{Char}(V_0) + \gamma_d(n, 1)\text{Char}(V_1) + \cdots + \gamma_d(n, d n)\text{Char}(V_{d n}).$$

We can summarize what we have shown so far in

Theorem 1.

$$\dim(C_d)_n = \omega_d\left(n, \frac{d n}{2}\right) + \omega_d\left(n, \frac{d n - 1}{2}\right).$$

Proof. The zero weight appears once in any representation V_k , for even k , therefore

$$\omega_d\left(n, \frac{d n}{2}\right) = \gamma_d(n, 0) + \gamma_d(n, 2) + \cdots + \gamma_d(n, 4) + \dots$$

The weight 1 appears once in any representation V_k , for odd k , therefore

$$\omega_d\left(n, \frac{d n - 1}{2}\right) = \gamma_d(n, 1) + \gamma_d(n, 3) + \cdots + \gamma_d(n, 5) + \dots$$

Thus

$$\omega_d\left(n, \frac{d n}{2}\right) + \omega_d\left(n, \frac{d n - 1}{2}\right) = \gamma_d(n, 0) + \gamma_d(n, 1) + \cdots + \gamma_d(n, d n).$$

□

3. It well-known that the number $\omega_d\left(n, \frac{d n}{2}\right)$ of non-negative integer solutions of the following system

$$\begin{cases} \alpha_1 + 2\alpha_2 + \cdots + d\alpha_d = \frac{d n}{2}, \\ \alpha_0 + \alpha_1 + \cdots + \alpha_d = n \end{cases}$$

is equal to the coefficient of $t^n z^{\frac{d n}{2}}$ of the expansion of the series

$$f_d(t, z) = \frac{1}{(1-t)(1-tz)\cdots(1-tz^d)}.$$

Denote it in such a way: $\omega_d\left(n, \frac{d n}{2}\right) = \left[t^n(z)^{\frac{d n}{2}}\right](f_d(t, z))$. It is clear that

$$\omega_d\left(n, \frac{d n}{2}\right) = [t^n z^{d n}](f_d(t, z^2)) = [(t z^d)^n](f_d(t, z^2)).$$

Similarly, the number $\omega_d\left(n, \frac{d n - 1}{2}\right)$ of non-negative integer solutions of the following system

$$\begin{cases} \alpha_1 + 2\alpha_2 + \cdots + d\alpha_d = \frac{d n - 1}{2}, \\ \alpha_0 + \alpha_1 + \cdots + \alpha_d = n \end{cases}$$

equals

$$\left[t^n(z)^{\frac{d n - 1}{2}}\right](f_d(t, z)) = [t^n z^{d n - 1}](f_d(t, z^2)) = [(t z^d)^n](z f_d(t, z^2)).$$

Thus, the following statement holds

Theorem 2. *The number $\dim(C_d)_n$ of linearly independent seminvariants (and covariants) of degree n for the binary d -form is calculated by the formula*

$$\dim(C_d)_n = [(t z^d)^n] \left(\frac{1 + z}{(1-t)(1-tz^2)\cdots(1-tz^{2d})} \right).$$

4. Let us prove Springer-type formula for the Poincaré series $P_d(z)$ of the algebra covariants of the binary d -form. Consider the $\mathbb{C}$ -algebra $\mathbb{C}[[t, z]]$ of formal power series. For arbitrary $m, n \in \mathbb{Z}^+$ define $\mathbb{C}$ -linear function

$$\Psi_{m,n} : \mathbb{C}[[t, z]] \rightarrow \mathbb{C}[[z]],$$

$m, n \in \mathbb{Z}^+$ in the following way

$$\Psi_{n_1, n_2}(t^{m_1} z^{m_2}) = \begin{cases} z^s, & \text{if } \frac{m_1}{n_1} = \frac{m_2}{n_2} = s \in \mathbb{N}, \\ 1, & \text{if } m_1 = m_2 = 0, \\ 0, & \text{otherwise.} \end{cases}$$

Then for arbitrary series

$$A = a_{0,0} + a_{1,0}t + a_{0,1}z + a_{2,0}t^2 + \dots,$$

we get

$$\Psi_{n_1, n_2}(A) = a_{0,0} + a_{n_1, n_2}z + a_{2n_1, 2n_2}z^2 + \dots.$$

Define by φ_n the restriction of $\Psi_{m,n}$ to $\mathbb{C}[[z]]$, namely

$$\varphi_n(z^m) := \begin{cases} z^{\frac{m}{n}}, & \text{if } m \equiv 0 \pmod{n}, \\ 0, & \text{if } m \not\equiv 0 \pmod{n}, \\ 1, & \text{for } m = 0. \end{cases}$$

It is clear that for arbitrary series

$$A = a_0 + a_1z + a_2z^2 + \dots,$$

we obtain

$$\varphi_n(A) = a_0 + a_nz + a_{2n}z^2 + \dots + a_{sn}z^s + \dots.$$

In some cases calculation of the functions Ψ can be reduced to calculation of the functions φ . The following statements hold:

Lema 2.

- (i) For $h(t, z) \in \mathbb{C}[[t, z]]$ we have $\Psi_{1,n}(h(t, z)) = \frac{1}{2\pi i} \oint_{|z|=1} h\left(\frac{t}{z^n}, z\right) \frac{dz}{z} \Big|_{t=z}$;
- (ii) for $R(z) \in \mathbb{C}[[z]]$ and for $m, n, k \in \mathbb{N}$ holds
$$\Psi_{1,n}\left(\frac{R(z)}{1 - tz^k}\right) = \begin{cases} \varphi_{n-k}(R(z)), & n \geq k, \\ 0, & \text{if } n < k \end{cases}$$

Proof. (i) Let $h(t, z) = \sum_{i,j=0}^{\infty} h_{i,j} t^i z^j$. Then

$$\frac{1}{2\pi i} \oint_{|z|=1} h\left(\frac{t}{z^n}, z\right) \frac{dz}{z} = \frac{1}{2\pi i} \sum_{i,j=0}^{\infty} \oint_{|z|=1} h_{i,j} t^i z^{j-ni} \frac{dz}{z} = \sum_i h_{i,ni} t^i.$$

Thus,

$$\frac{1}{2\pi i} \oint_{|z|=1} h\left(\frac{t}{z^n}, z\right) \frac{dz}{z} \Big|_{t=z} = \sum_i h_{i,ni} z^i = \Psi_{1,n}(h(t, z)).$$

(ii) Let $R(z) = \sum_{j=0}^{\infty} f_j z^j$. Then for $k < n$ we have

$$\Psi_{m,n}\left(\frac{R(z)}{1 - t^m z^k}\right) = \Psi_{m,n}\left(\sum_{j,s=0}^{\infty} f_j z^j (t^m z^k)^s\right) = \Psi_{m,n}\left(\sum_{s=0}^{\infty} f_{s(n-k)} (t^m z^n)^s\right) = \sum_{s=0}^{\infty} f_{s(n-k)} z^s.$$

On other hand, $\varphi_{n-k}(R(z)) = \varphi_{n-k}\left(\sum_{j=0}^{\infty} f_j z^j\right) = \sum_{s \geq 0} f_{s(n-k)} z^s$.

□

The main idea of this calculations is that the Poincaré series $P_d(z)$ can be expressed in terms of functions Ψ . The following simple but important statement holds

Lema 3.

$$P_d(z) = \Psi_{1,d} \left(\frac{1+z}{(1-t)(1-tz^2)\dots(1-tz^{2d})} \right).$$

Proof. Theorem 2 implies that $\dim(C_d)_n = [(tz^d)^n]f_d(t, z^2)$. Then

$$P_d(z) = \sum_{n=0}^{\infty} \dim(C_{2,d})_n z^n = \sum_{n=0}^{\infty} ([(tz^d)^n]f_d(t, z^2)) z^n = \Psi_{1,d}(f_d(t, z^2)).$$

□

Combining this with Lemma 2, (i) we obtain one more formula for the Puanaré series:

$$P_d(t) = \frac{1}{2\pi i} \oint_{|r|=1} \frac{1+z}{\prod_{k=0}^d (1-tz^{d-2k})} \frac{dz}{z}.$$

Compare the formula with the Molien-Weyl integral formula for the Poincaré series of the algebra of invariants of binary form, see [8], p. 183.

Now we can present Springer-type formula for the Poincaré series $P_d(z)$

Theorem 3.

$$P_d(z) = \sum_{0 \leq k < d/2} \varphi_{d-2k} \left(\frac{(-1)^k z^{k(k+1)} (1+z)}{(z^2, z^2)_k (z^2, z^2)_{d-k}} \right),$$

here $(a, q)_n = (1-a)(1-aq)\dots(1-aq^{n-1})$ is q -shifted factorial.

Proof. Consider the partial fraction decomposition of the rational function $f_d(t, z^2)$

$$f_d(t, z^2) = \sum_{k=0}^d \frac{R_k(z)}{1-tz^{2k}}.$$

It is easy to see, that

$$\begin{aligned} R_k(z) &= \lim_{t \rightarrow z^{-2k}} (f_d(t, z^2)(1-tz^k)) = \lim_{t \rightarrow z^{-2k}} \left(\frac{(1+z)}{(t, z)_{d+1}} (1-tz^{2k}) \right) = \\ &= \frac{1+z}{(1-z^{-2k})(1-z^{2-2k})\dots(1-z^{2(k-1)-2k})(1-z^{2(k+1)-2k})\dots(1-z^{2d-2k})} = \\ &= \frac{z^{2k+(2k-2)+\dots+2}(1+z)}{(z^{2k}-1)(z^{2k-2}-1)\dots(z^2-1)(1-z^2)\dots(1-z^{2d-2k})} = \frac{(-1)^k z^{k(k+1)} (1+z)}{(z^2, z^2)_k (z^2, z^2)_{d-k}}. \end{aligned}$$

Using the above lemmas we obtain

$$P_d(z) = \Psi_{1,d}(f_d(t, z^2)) = \Psi_{1,d} \left(\sum_{k=0}^n \frac{R_k(z^2)}{1-tz^{2k}} \right) = \sum_{0 \leq k < d/2} \varphi_{d-2k} \left(\frac{(-1)^k z^{k(k+1)} (1+z)}{(z^2, z^2)_k (z^2, z^2)_{d-k}} \right).$$

□

5. For direct computations we use the following technical lemma

Lema 4. *Let $R(z)$ be some polynomial of z . Then*

$$\varphi_n \left(\frac{R(z)}{(1-z^{k_1})(1-z^{k_2}) \cdots (1-z^{k_m})} \right) = \frac{\varphi_n(R(z)Q_n(z^{k_1})Q_n(z^{k_2})Q_n(z^{k_m}))}{(1-z^{k_1})(1-z^{k_2}) \cdots (1-z^{k_m})},$$

here $Q_n(z) = 1 + z + z^2 + \cdots + z^{n-1}$, and k_i are natural numbers.

Proof. Observe that $\varphi_n(f(z^n)g(z)) = f(z)\varphi_n(g(z))$, for arbitrary $f(z), g(z) \in \mathbb{C}[[z]]$. Particularly,

$$\begin{aligned} \varphi_n \left(\frac{g(z)}{1-z^m} \right) &= \varphi_n \left(\frac{R(z)}{1-z^{nm}} \frac{1-z^{nm}}{1-z^m} \right) = \frac{1}{1-z^m} \varphi_n \left(g(z) \frac{1-z^{nm}}{1-z^m} \right) = \\ &= \frac{1}{1-z^m} \varphi_n (g(z)(1+z^m+(z^m)^2+\cdots+(z^m)^{n-1})) = \frac{1}{1-z^m} \varphi_n (g(z)Q_n(z^m)). \end{aligned}$$

□

Example. Consider the case $d = 3$. We have

$$\begin{aligned} P_3(z) &= \Psi_{1,3} \left(\frac{1+z}{(1-t)(1-tz^2)(1-tz^4)(1-tz^6)} \right) = \\ &= \varphi_3 \left(\frac{1+z}{(1-z^2)(1-z^4)(1-z^6)} \right) - \varphi_1 \left(\frac{(1+z)z^2}{(1-z^2)^2(1-z^4)} \right) = \\ &= \frac{1}{1-z^2} \varphi_3 \left(\frac{1+z}{(1-z^6)(1-z^{12})} \cdot \frac{1-z^6}{1-z^2} \cdot \frac{1-z^{12}}{1-z^4} \right) - \frac{(1+z)z^2}{(1-z^2)^2(1-z^4)} = \\ &= \frac{1}{(1-z^2)^2(1-z^4)} \varphi_3 ((1+z)(1+z^2+z^4)(1+z^4+z^8)) - \frac{(1+z)z^2}{(1-z^2)^2(1-z^4)} = \\ &= \frac{1+z+z^2+2z^3+z^4}{(1-z^2)^2(1-z^4)} - \frac{(1+z)z^2}{(1-z^2)^2(1-z^4)} = \frac{1+z+z^3+z^4}{(1-z^2)^2(1-z^4)} = \\ &= \frac{(1+z)(1+z^3)}{(1-z^2)^2(1-z^4)} = \frac{1+z^3}{(1-z)(1-z^2)(1-z^4)}. \end{aligned}$$

By using Lemma 3 the Poncaré series $P_d(z)$ for $d \leq 20$ are found. Below list several results:

$$\begin{aligned} P_1(z) &:= \frac{1}{1-z}, P_2(z) := \frac{1}{(-1+z^2)(z-1)}, \\ P_3(z) &:= \frac{1+z^3}{(1-z)(1-z^2)(1-z^4)}, P_4(z) := \frac{1+z^3}{(1-z)(1-z^2)^2(1-z^3)} \\ P_5(z) &:= \frac{z^{15}+z^{13}+3z^{12}+3z^{11}+5z^{10}+4z^9+6z^8+6z^7+4z^6+5z^5+3z^4+3z^3+z^2+1}{(1-z)(1-z^2)(1-z^4)(1-z^6)(1-z^8)} \\ P_6(z) &= \frac{z^{10}+z^8+3z^7+4z^6+4z^5+4z^4+3z^3+z^2+1}{(1-z)(1-z^2)^2(1-z^3)(1-z^4)(1-z^5)} \\ P_7(z) &:= p_7(z)/((1-z)(1-z^2)(1-z^4)(1-z^6)(1-z^8)(1-z^{10})(1-z^{12})) \end{aligned}$$

where

$$\begin{aligned} p_7(z) := & z^{35} + 2z^{33} + 6z^{32} + 10z^{31} + 19z^{30} + 28z^{29} + 44z^{28} + 61z^{27} + 79z^{26} + 102z^{25} + \\ & + 129z^{24} + 156z^{23} + 173z^{22} + 196z^{21} + 215z^{20} + 230z^{19} + 231z^{18} + 231z^{17} + \\ & + 230z^{16} + 215z^{15} + 196z^{14} + 173z^{13} + 156z^{12} + 129z^{11} + 102z^{10} + 79z^9 + \\ & + 61z^8 + 44z^7 + 28z^6 + 19z^5 + 10z^4 + 6z^3 + 2z^2 + 1. \end{aligned}$$

$$P_8(z) = p_8(z)/((1-z)(1-z^2)^2(1-z^3)^2(1-z^4)(1-z^5)(1-z^7))$$

$$\begin{aligned} p_8(z) = & z^{18} + 2z^{16} + 6z^{15} + 12z^{14} + 19z^{13} + 25z^{12} + 31z^{11} + 36z^{10} + 38z^9 + 36z^8 + \\ & + 31z^7 + 25z^6 + 19z^5 + 12z^4 + 6z^3 + 2z^2 + 1 \end{aligned}$$

$$P_9(z) = p_9(z)/(1-z)(1-z^2)(1-z^4)(1-z^6)(1-z^8)(1-z^{10})(1-z^{12})(1-z^{14})(1-z^{16}),$$

$$\begin{aligned} p_9(z) = & z^{63} + 3z^{61} + 10z^{60} + 23z^{59} + 49z^{58} + 93z^{57} + 172z^{56} + 289z^{55} + 457z^{54} + 701z^{53} + \\ & + 1036z^{52} + 1477z^{51} + 2023z^{50} + 2720z^{49} + 3568z^{48} + 4573z^{47} + 5702z^{46} + \\ & + 7013z^{45} + 8466z^{44} + 10043z^{43} + 11672z^{42} + 13400z^{41} + 15155z^{40} + 16880z^{39} + \\ & + 18487z^{38} + 20013z^{37} + 21392z^{36} + 22539z^{35} + 23398z^{34} + 24013z^{33} + \\ & + 24355z^{32} + 24355z^{31} + 24013z^{30} + 23398z^{29} + 22539z^{28} + 21392z^{27} + \\ & + 20013z^{26} + 18487z^{25} + 16880z^{24} + 15155z^{23} + 13400z^{22} + 11672z^{21} + \\ & + 10043z^{20} + 8466z^{19} + 7013z^{18} + 5702z^{17} + 4573z^{16} + 3568z^{15} + 2720z^{14} + \\ & + 2023z^{13} + 1477z^{12} + 1036z^{11} + 701z^{10} + 457z^9 + 289z^8 + 172z^7 + 93z^6 + \\ & + 49z^5 + 23z^4 + 10z^3 + 3z^2 + 1 \end{aligned}$$

$$P_{10}(z) = p_{10}(z)/(1-z)(1-z^2)^2(1-z^3)(1-z^4)(1-z^5)(1-z^6)(1-z^7)(1-z^8)(1-z^9),$$

$$\begin{aligned} p_{10}(z) = & z^{36} + 3z^{34} + 11z^{33} + 27z^{32} + 58z^{31} + 112z^{30} + 193z^{29} + 318z^{28} + 485z^{27} + 699z^{26} + \\ & + 951z^{25} + 1245z^{24} + 1541z^{23} + 1842z^{22} + 2108z^{21} + 2321z^{20} + 2451z^{19} + \\ & + 2506z^{18} + 2451z^{17} + 2321z^{16} + 2108z^{15} + 1842z^{14} + 1541z^{13} + 1245z^{12} + \\ & + 951z^{11} + 699z^{10} + 485z^9 + 318z^8 + 193z^7 + 112z^6 + 58z^5 + 27z^4 + 11z^3 + \\ & + 3z^2 + 1. \end{aligned}$$

REFERENCES

- [1] Sylvester, J. J., Franklin, F., Tables of the generating functions and groundforms for the binary quantic of the first ten orders, Am. J. II., 223–251, 1879
- [2] Sylvester, J. J. Tables of the generating functions and groundforms of the binary duodecimic, with some general remarks, and tables of the irreducible syzigies of certain quantics. Am. J. IV. 41–62, 1881.
- [3] Springer T.A., On the invariant theory of $SU(2)$, Indag. Math. 42 (1980), 339–345.
- [4] Brouwer A., Cohen A., The Poincaré series of the polynomial invariants under SU_2 in its irreducible representation of degree ≤ 17 , preprint of the Mathematisch Centrum, Amsterdam, 1979.
- [5] Littelman P., Procesi C., On the Poincaré series of the invariants of binary forms, J. Algebra 133, No.2, 490–499 (1990).
- [6] Roberts M., The covariants of a binary quantic of the n -th degree, Quarterly J. Math., 4, 1861, 168–178.
- [7] W. Fulton, J. Harris, Representation theory: a first course, 1991.
- [8] Derksen H., Kemper G., Computational Invariant Theory, Springer-Verlag, New York, 2002.

KHMELNITSKIY NATIONAL UNIVERSITY, INSITUTS'KA, 11, KHMELNITSKIY, 29016, UKRAINE